


**Chemistry Atomic Structure and Chemical Bonding**  
**Prof. K. Mangala Sunder**  
**Department of Chemistry**  
**Indian Institute of Technology, Madras**

**Lecture - 09**  
**Particle in a two dimensional box (Infinite Barrier)**

Welcome back to the lecture the earlier lecture talked about in the earlier lecture I talked about the particle in a one dimensional box and in the current one lets discuss the particle in a two dimensional two dimensional model or two degrees of freedom model. The particles position coordinates are given by two x and y, two coordinates in a plane. Orthogonal to each other and then we discussed the quantum problem

(Refer Slide Time: 00:52)




The barriers are infinite therefore if you remember the problem  $p^2$  by  $2m$  plus  $v$  which is the energy term gets changed to or its re written as  $p_x^2$  by  $2m$  plus  $p_y^2$  by  $2m$  plus  $v$  and  $p_x$  is replaced in quantum mechanics by the minus  $\hbar$  square by the term minus  $\hbar$  square by  $2m$  the partial derivative now.

Because we have the wave function as a function of two coordinates  $x$  and  $y$  and the momentum in the  $x$  direction is given by the partial derivative and this is the square of the momentum. So, you have minus  $\hbar$  square  $\frac{\partial^2}{\partial x^2}$  by  $2m$  and correspondingly for  $p_y$  squared you have  $\frac{\partial^2}{\partial y^2}$  this is the operator

part for the kinetic energy of the Hamiltonian plus and the wave function is a function of  $x$  and  $y$  plus  $V$  some potential times  $\psi(x, y)$  is equal to  $E \psi(x, y)$ .

This is the two dimensional Schrodinger equation in which you have got the  $\hbar$  this term plus the  $\nabla^2$  acting on the  $\psi$  giving you  $E \psi$  and for the current problem of particle in the two d box. We considered  $V$  to be infinite for all values of  $x$  other than from 0 to  $L$  and all values of  $y$  from 0 to some other say  $a$  or  $L_1$  or  $L_2$  it does not matter if it is a rectangular box.


(Refer Slide Time: 03:03)



If it is a square box, then essentially you are looking at the not see if you can have a square something like that. So, 0 to  $L$  and  $y$  is also 0 to  $L$  only in this region we are looking at the particle properties and the particles behavior and for all others we have  $V$  is infinity for all values of  $x$  less than 0 or equal to and for all values of  $x$  greater than or equal to  $L$  and likewise for  $y$  less than or equal to 0  $y$  greater than or equal to  $L$ . So, this is the infinite boundaries that you have its not the single dimensional quantity, but it is the surface in a sense that we protect the particle from escaping this region and inside  $V$  is 0 between  $x$  and  $L$  between  $y$  and  $L$  and this is a square box.

So, if we do that; obviously, the differential equation simplifies without this term and you have a derivative square in one direction a derivative square in another direction and then you have the  $\psi$  of  $x, y$  ok.


(Refer Slide Time: 04:49)



Such a problem is easily solved by is written in terms of a product of a function of x alone and a function of y alone ok. With this choice it is possible to separate this equation minus h bar square by 2m dou square by dou x square plus dou square by dou y square psi of x comma y is equal to E times psi of x comma y into two equations namely minus h bar square by 2m d square by d x square x is equal to E 1 of x and minus h bar square by 2m d square by d y square times y is equal to E 2 times y, but these two constants E 1 and E 2 are constrained by E 1 plus E 2 is equal to E ok.


The actual separation of this is given in the notes that accompanies this video lecture. Therefore, I would request you to look into that to see how this equation is separated into two one dimensional equation one for x and one for y with the constraint that the energies for the two one dimensional problems are related to the total energy as the sum E 1 plus E 2.

(Refer Slide Time: 06:49)



independently of the other. Therefore, the above equation is satisfied only if each term is separately equal to a constant.

$$-\frac{1}{X(x)} \frac{\hbar^2}{2m} \frac{d^2 X(x)}{dx^2} = E_1, \text{ or, } -\frac{\hbar^2}{2m} \frac{d^2 X(x)}{dx^2} = E_1 X(x) \text{ and}$$
$$-\frac{1}{Y(y)} \frac{\hbar^2}{2m} \frac{d^2 Y(y)}{dy^2} = E_2, \text{ or, } -\frac{\hbar^2}{2m} \frac{d^2 Y(y)}{dy^2} = E_2 Y(y)$$



Now let us see the solutions that quantity which I have written on the board is namely this is the x equation and the corresponding y equation is that ok. Obviously, each one of them is like a one dimensional part particle in a box.

Therefore the solutions for each one of them will have a running quantum number for that particular equation. The x component of the wave function will be given by the solution its similar to the psi of x that we wrote, except that now we call it x of x and now this will have a quantum number going from 1,2,3 to some value which we call as n1. In an exactly in an identical manner the y equation will also have a free quantum number n2 which will run from 1,2,3 to whatever that we take.

But please remember these two quantum numbers are not independent in the sense they are connected to the total energy the requirement that E1 plus E2 is equal to E.

(Refer Slide Time: 08:18)

$\psi(x, y)$

$E_1 = \frac{h^2}{8mL^2} n_1^2$

$E_2 = \frac{h^2}{8mL^2} n_2^2$

$E_1$  and  $E_2$  satisfy the condition

$E_1 + E_2 = E$

Separation of variables

$\left. \begin{array}{l} E_1 = \frac{h^2}{8mL^2} n_1^2 \\ E_2 = \frac{h^2}{8mL^2} n_2^2 \end{array} \right\} \frac{h^2}{8mL^2} (n_1^2 + n_2^2) = E$

What are the new results for the particle in the two dimensional (coordinate) square box whose side is of length L. Then we have all the solutions same as one dimensional box except that they are repeated for one coordinate each.

dimensional problem. An important new result in addition is the requirement that the total energy is the sum of the two one dimensional energies.

NPTEL

Now remember the expression for  $E_1$  from the particle in a one dimensional box it is  $\frac{h^2}{8mL^2} n_1^2$  a free quantum number in the sense it takes 1, 2, 3 integer values and  $E_2$  is also given by  $\frac{h^2}{8mL^2} n_2^2$  such that this equation is satisfied. Therefore, you have  $\frac{h^2}{8mL^2} n_1^2 + \frac{h^2}{8mL^2} n_2^2 = E$ .

So, this is the only constraint that comes out in the separation of the two dimensional Schrödinger equation that the total energy is the sum of the two one dimensional energies and that is possible because we do not have a potential which couples the two dimensions, we have put  $V$  is equal to zero and therefore, the method of separation of variables. Separation of variables we have separated the  $x$  and  $y$  from the  $\psi$  of  $x, y$  if you recall. The  $\psi$  of  $x, y$  they have separated that into the  $x$  equation and the  $y$  equation. So, that process is called the separation of variables now how do these functions look like.

(Refer Slide Time: 10:06)

dimensional problem. An important new result in addition to the require-  
energies have to add up to a total energy, which will lead to the idea of  
states, to be discussed below.

$$X_{n_1}(x) = \sqrt{\frac{2}{L}} \sin\left(\frac{n_1 \pi x}{L}\right), E_{n_1} = \frac{h^2 n_1^2}{8mL^2}$$

$$Y_{n_2}(y) = \sqrt{\frac{2}{L}} \sin\left(\frac{n_2 \pi y}{L}\right), E_{n_2} = \frac{h^2 n_2^2}{8mL^2}$$

$$E_{n_1} + E_{n_2} = \frac{h^2}{8mL^2} (n_1^2 + n_2^2) = E_{n_1 n_2}$$

Obviously you have the solutions for the quantum number  $n_1$  in terms of the one dimensional solution that you have seen in the previous lecture. Root 2 by L sin  $n_1 \pi x$  by l and the energy is given by  $n_1$  square and likewise for the y with the  $n_2$  square and with the constraint that the total energy  $E_{n_1} + E_{n_2}$  is  $E_{n_1 n_2}$  you have seen that ok.

(Refer Slide Time: 10:43)

$$Y_{n_2}(y) = \sqrt{\frac{2}{L}} \sin\left(\frac{n_2 \pi y}{L}\right), E_{n_2} = \frac{h^2 n_2^2}{8mL^2}$$

$$E_{n_1} + E_{n_2} = \frac{h^2}{8mL^2} (n_1^2 + n_2^2) = E_{n_1 n_2}$$

$$\Psi_{n_1 n_2}(x, y) = X_{n_1}(x) Y_{n_2}(y)$$

Examples of quantum numbers, wave functions and energies:

$n_1 = 1, n_2 = 1$	$\psi_{11}(x, y) = \frac{2}{L} \sin\left(\frac{\pi x}{L}\right) \sin\left(\frac{\pi y}{L}\right)$	$E_{11} = \frac{h^2}{8mL^2} (1^2 + 1^2)$
--------------------	---	--

What about the wave function? The wave function now if you see this the wave function  $\psi$  of  $n_1 n_2$  because it is; obviously, specified by the two quantum numbers  $n_1$  and  $n_2$

has the independent function x with the quantum number n 1 and y with the quantum number n 2 each one is in an orthogonal direction ok.

Therefore you see this interesting thing next line, when we have n 1 is 1 and n 2 is 1 when we have that case which is the starting point what is called the lowest energy for the particle in a two dimensional box. You can see that the wave function is given by psi 1 1 x comma y and is given by the product of the two functions that you saw the x of x and y of y which gives you sin pi x by L and sin pi y by L.

Let me repeat this when the quantum number is 1 1 the wave function is given by psi 1 1 and its given by the product of 2 by L sin pi x by L and sin pi y by L and the energy is of course, the sum of one square plus one square times the whole thing. Therefore, the energy for this process E 1 1 is h square by 8m l square times 2.

(Refer Slide Time: 12:06)

The screenshot shows a presentation slide with the following content:

- Equation:  $\left(\frac{n_2 \pi y}{L}\right), E_{n_2} = \frac{h^2 n_2^2}{8mL^2}$
- Equation:  $(n_1^2 + n_2^2) = E_{n_1, n_2}$
- Equation:  $(\cdot) Y_{n_2}(y)$
- Text: "um numbers, wave functions and energies:"
- Equation:  $\psi_{n_1, n_2}(x, y) = \frac{2}{L} \sin\left(\frac{\pi x}{L}\right) \sin\left(\frac{\pi y}{L}\right)$
- Equation:  $E_{11} = \frac{h^2}{8mL^2} (1^2 + 1^2) = \frac{h^2}{4mL^2}$
- Handwritten notes:
  - $\psi_{n_1, n_2} = X_{n_1} Y_{n_2}$  with  $n_1 \neq n_2$
  - $= X_{n_2} Y_{n_1}$  with  $E \rightarrow (n_1^2 + n_2^2) \frac{h^2}{8mL^2}$
  - "Degenerate state (2)"

What is interesting is the next choice you have psi n 1 n 2 as x of n 1 y of n2. Its possible if n 1 is not equal to n 2 it is possible to have the wave function given by x of n 2 and y of n 1 because the energy is simply proportional to n 1 square plus n 2 square.

Times h square by of course, 8m L square which is the proportionality constant. Therefore, you see that you have the same energy, but you have two physically different states x of n 1 y of n 2 and x of n 2 y of n 1 both states have the same energy this is what is called a degenerate state.

Degeneracy is 2 because there are two states which have the same quantum same energy, but they have different quantum states. This is the introduction for the particle in a two d box that the degeneracy is the additional factor. Now how do these things look like let us simplify this picture. Now I have a whole series of functions here with which you can fill up any number of pages if you wish.

(Refer Slide Time: 13:51)

Examples of quantum numbers, wave functions and energies:

$n_1 = 1, n_2 = 1$	$\psi_{11}(x, y) = \frac{2}{L} \sin\left(\frac{\pi x}{L}\right) \sin\left(\frac{\pi y}{L}\right)$	$E_{11} = \frac{\hbar^2}{8mL^2}(1^2 + 1^2) = \frac{\hbar^2}{4mL^2}$
$n_1 = 2, n_2 = 1$	$\psi_{21}(x, y) = \frac{2}{L} \sin\left(\frac{2\pi x}{L}\right) \sin\left(\frac{\pi y}{L}\right)$	$E_{21} = \frac{\hbar^2}{8mL^2}(2^2 + 1^2) = \frac{5\hbar^2}{8mL^2}$
$n_1 = 1, n_2 = 2$	$\psi_{12}(x, y) = \frac{2}{L} \sin\left(\frac{\pi x}{L}\right) \sin\left(\frac{2\pi y}{L}\right)$	$E_{12} = \frac{\hbar^2}{8mL^2}(1^2 + 2^2) = \frac{5\hbar^2}{8mL^2}$
$n_1 = 2, n_2 = 2$	$\psi_{22}(x, y) = \frac{2}{L} \sin\left(\frac{2\pi x}{L}\right) \sin\left(\frac{2\pi y}{L}\right)$	$E_{22} = \frac{\hbar^2}{8mL^2}(2^2 + 2^2) = \frac{\hbar^2}{mL^2}$
$n_1 = 3, n_2 = 2$	$\psi_{32}(x, y) = \frac{2}{L} \sin\left(\frac{3\pi x}{L}\right) \sin\left(\frac{2\pi y}{L}\right)$	$E_{32} = \frac{\hbar^2}{8mL^2}(3^2 + 2^2) = \frac{13\hbar^2}{8mL^2}$

Handwritten notes:  $E_{n_1, n_2} = \frac{\hbar^2}{8mL^2}(n_1^2 + n_2^2) = E_{n_2, n_1}$   
 $\Psi_{n_1, n_2}(x, y) = X_{n_1}(x)Y_{n_2}(y)$   
 $E \rightarrow (n_1^2 + n_2^2) \frac{\hbar^2}{8mL^2}$   
 Degenerate state (2)

You see that  $n_1 = 2, n_2 = 1$  corresponds to the wave function  $\psi_{21}$  with  $\sin 2\pi x \sin \pi y$  by  $L$ , and  $n_1 = 1, n_2 = 2$  gives you the other function namely  $\sin \pi x \sin 2\pi y$  by  $L$  and the energies are the same.

So, if the quantum numbers are identical there is no degeneracy, but if the quantum numbers are different for a square box because we have chosen the length  $L$  to be the same the square box gives you the solution that you have a minimum degeneracy of 2 if  $n_1$  is not the same as  $n_2$  and you can see that for 3 and 2 that you have here the wave function  $\sin 3\pi x$  by  $L$  and  $\sin 2\pi y$  by  $L$  and then 2 and 3 which is  $\sin 2\pi x$  by  $L$  and  $\sin 3\pi y$  by  $L$ .



(Refer Slide Time: 14:46)

iii) confined to move in a square box whose length is  $L$  and whose all sides have infinite

$n_1 = 3, n_2 = 2$	$\psi_{32}(x,y) = \frac{2}{L} \sin\left(\frac{3\pi x}{L}\right) \sin\left(\frac{2\pi y}{L}\right)$	$E_{12} = \frac{\hbar^2}{8mL^2}(3^2 + 2^2) = \frac{13\hbar^2}{8mL^2}$
$n_1 = 2, n_2 = 3$	$\psi_{23}(x,y) = \frac{2}{L} \sin\left(\frac{2\pi x}{L}\right) \sin\left(\frac{3\pi y}{L}\right)$	$E_{23} = \frac{\hbar^2}{8mL^2}(2^2 + 3^2) = \frac{13\hbar^2}{8mL^2}$

Thus,

- the particle not having any potential energy inside a square region,
- moving in a two dimensional plane,
- confined to move in a square box whose length is  $L$  and whose all sides have infinite

So, the axes choice the quantum number choice for a given axes determines the functions state.

(Refer Slide Time: 15:06)

iii) confined to move in a square box whose length is  $L$  and whose all sides have infinite repulsive potential and

iv) obeying the Schrodinger equation has its energy quantized in terms of two quantum numbers  $n_1$  and  $n_2$  such that they determine the total energy of the particle which is now discrete and can have multiple states having the same energy.

What is new besides an additional quantum number?

ANSWER: DEGENERACY (Same energy for more than one possible state of the particle)


$\Psi_{n_1 n_2}(x,y) = X_{n_1}(x)Y_{n_2}(y)$
$\Psi_{n_2 n_1}(x,y) = X_{n_2}(x)Y_{n_1}(y)$

Two questions for you. Are they two different states of the particle? What about their energy?

Degenerate energy levels in units of  $\frac{\hbar^2}{8mL^2}$  of a particle in a two-d box energy levels and three dimensional graphical representation of a few wave functions and their absolute squares are included in the lecture.


How do these things look like if we plot them, I mean this plot looks fancy, but actually does not have much interpretation or meaning.

(Refer Slide Time: 15:08)




But it is what seeing the product wave function in two dimensions.

(Refer Slide Time: 15:11)




So, you see the wave function yeah you see the wave function  $\psi_{11}$  using this picture. It is a half wave similar to what you had in your particle in a one dimensional box in the x direction and it is also a half wave in the y direction as you can see through the projection, in the x direction here of this graph and on the y direction also you have the same thing identical ok.

(Refer Slide Time: 15:56)



What about the sin square?

(Refer Slide Time: 16:03)




The sin square which is associated with the probability that the particle be found not in a small length region  $dx$ , but in a small area  $dx dy$  please remember  $\psi \times y$  if you do that  $\psi$  square  $dx dy$  is the probability that the particle will be in the small rectangular region between  $x$  and  $x$  plus  $dx$  and  $y$  and  $y$  plus  $dy$ .

That is a small region and you can see that the  $\psi$  square is given like this therefore, you can create I mean you can visualize what would be the probability exactly the same way

that you have visualized the particle in a one dimensional box, except that now we have a motion on the plane and now what is interesting is, when you go to different quantum numbers where there is degeneracy.

(Refer Slide Time: 17:11)



Psi 1 2 if you look at this psi 1 2 is quantum number 1 for the x direction and quantum number 2 for the y direction.


This is the quantum number for the x direction and you can see that it is a half wave which is either up or down it is either positive or negative. The reason being the y direction wave is a full wave. So, in this direction what you have is if I may draw this the wave function looks like that, in this direction the wave function looks like that. Therefore, when you take the product of these two functions a negative side makes this wave function negative for half the length and therefore, you see that for half the length you have either a positive wave function or you have a negative wave function.

That is only for the wave function, we know that the wave function is not that important it is the square of the wave function which is important for probability interpretation and you can see that psi square which removes this negative character of the function gives you now very beautifully the 2 n equal to 1 case for the x axis and the n equal to 2. If you remember the graph that you had for n equal to 2 for the y axis and this is the x axis.

Therefore the features are captured the wave function features are captured when you do your surface plot and you can see that the pictures can be created for a large number of them, but there is element two dimensions and in three dimension we probably can use colour at the most to distinguish the function from the three axis, but that is it you cannot visualize this for n dimensions.

So, let us conclude this part of the particle in a two dimensional box with some examples of the wave functions and the squares of the wave function for different quantum numbers.

(Refer Slide Time: 19:25)




So, here is a 2 1 as opposed to 1 2 and you see all that happens is that for a 2 1. the wave function along the x axis is like this and the wave function along the y axis it is like that and you can see that actually sorry this is in the wrong direction so let me erase that because your 0 starts from here.

Therefore, you have that and this is the y axis that is the reason why part of it is negative and the other part is positive and the square of the wave function you can see that there are two humps along the x axis and along the y axis it is a quantum number 1. So, you have only 1.

Similar to the one dimensional y axis and let us see 1 or 2 more examples and let me stop with that.

(Refer Slide Time: 20:29)



This is a I mean the exercise here what does this picture represent? There is 1 here along the x axis and there are 3 peaks therefore, you have this is a y is 3 and x is 1. So, it is psi 1 3 squared x y.

So, the lecture notes give you many more such pictures, but in the next part of this lecture we will see what do all these things mean in terms of probability calculations and in terms of a new idea called the expectation values. We will stop here for this particular part of the lecture.

Thank you.